


# New definitions of scientific units are on the horizon

Metrologists are poised to change how scientists measure the Universe.

Elizabeth Gibney

18 October 2017



A sphere of pure silicon can be used to define a unit of measurement known as a mole.

Natl. Phys. Lab., UK

Revamped definitions of scientific units are on their way. In the biggest overhaul of the international system of units (SI) since its inception in 1960, a committee is set to redefine four basic units — the ampere, the kilogram, the kelvin and the mole — using relationships to fundamental constants, rather than abstract or arbitrary definitions. The International Bureau of Weights and Measures is reviewing the plans at a meeting near Paris from 16 to 20 October. Its recommendations will then go before the General Conference on Weights and Measures, which oversees the SI system, in November 2018. The changes would take effect in May 2019.

The kilogram is currently defined as the mass of a chunk of metal in a vault in Paris. And an imaginary experiment involving the force between two infinite wires defines the ampere, the unit of electrical current. The mole, meanwhile, is the amount of substance in a system with as many elementary entities as there are atoms in 0.012 kilograms of carbon-12, while the kelvin relates to the temperature and pressure at which water, ice and water vapour co-exist in equilibrium, known as the triple point of water. In the future, these units will be calculated in relation to constants — for example, the ampere will be based on the charge of an electron.

Redefinition might not affect everyday measurements, but it will enable scientists working at the highest level of precision to do so in multiple ways, at any place or time and on any scale, without losing accuracy.


**Related stories**

- The new system of units
- Kilogram conflict resolved at last
- Hyper-precise atomic clocks face off to redefine time

**More related stories**

## ALL CHANGE

Under the revised SI system, every unit will be defined in relation to a constant, whose value will become fixed. Many of the units will be defined in relation to each other; for example, definition of the kilogram requires Planck's constant, and definitions of the second and metre.\*




**The problem**

For measurements on conventional scales, existing definitions of SI units suffice. But they are poor tools for modern science at the extremes. And basing units on specific points or materials can be troublesome and inelegant, say metrologists.


## THE UNSTABLE KILOGRAM

The kilogram is currently defined by a lump of platinum-iridium, stored in a vault near Paris. Because objects can easily lose atoms or absorb molecules from the air, using one to define an SI unit is problematic. Compared to the prototype, some official copies have gained at least 50 micrograms over a century.



## A QUESTION OF SCALE

When a unit is defined on a fixed scale, uncertainties grow larger the further scientists move away from that point. Currently, for example, measurements in milligram have a minimum relative uncertainty 2,500 times that associated with the kilogram. The problem disappears under the proposed system, which relies on constants to define units.




Source: Shaw, G. et al. Metrologia 53, A86-A94 (2016).

**The techniques**

Under the revamped SI system, researchers will be able to use various experiments to relate constants to each of the units measured.


## AMPERE: THE SINGLE-ELECTRON PUMP

Used to measure the charge of an electron, an electron pump could become one tool for determining the ampere. By trapping individual electrons as they travel rapidly across a conductor, the pump can generate a measurable current by counting single electrons.




## MOLE: THE SILICON SPHERE

As the device that gives scientists Avogadro's constant, this silicon sphere offers a state-of-the-art way to measure a mole. It would determine the precise number of atoms in a perfect sphere of pure silicon-28. Researchers do this by using lasers to measure the length of a unit of the sphere's crystal lattice, and its mean diameter.




## KILOGRAM: THE WATT BALANCE

The Watt balance compares mechanical power with electromagnetic power using two separate experiments. First, a current is run through a coil in a magnetic field to create a force that counterbalances a known physical mass. Then, the coil is moved through the field to create a voltage. By measuring the speed as well as experimental values that relate the voltage and current to Planck's constant, scientists can precisely determine the weight of a mass in kilograms.



## KELVIN: ACOUSTIC THERMOMETRY


This technique could be used to derive precise temperature measurements. The speed of sound in a gas-filled sphere (which is proportional to the average speed of the atoms in it) can be determined at a fixed temperature, by analysing the frequency of sound waves that resonate within it and measuring the sphere's volume.



## THE FUTURE

Experimental teams have been working for decades to agree on values for the constants on which the definitions will soon hinge. They had to meet strict conditions, which the kilogram teams fulfilled only in 2015. All groups submitted final figures by 1 July. Under the new system, these constants will be stripped of their uncertainties and fixed as exact numbers in May 2019. Their former uncertainties will then be transferred to measurements that use the units defined by the constants. As a consequence, other, related constants, once expressed in the new units, will see their uncertainties reduced as well.

The loser will be the mass of the prototype kilogram in Paris. It currently has an uncertainty of zero — but that will soon rise to at least ten parts per billion.



Nature 550, 312–313 (19 October 2017) doi:10.1038/550312a

**Related stories and links**

**From nature.com**

- The new system of units 07 January 2016
- Kilogram conflict resolved at last 14 October 2015
- Hyper-precise atomic clocks face off to redefine time 02 June 2015
- Gravity rivals join forces to nail down Big G 08 October 2014
- Ampere to get rational redefinition 14 January 2014
- Metrology: The new and improved kelvin 17 June 2009

**From elsewhere**

- International Bureau of Weights and Measures

For the best commenting experience, please login or register as a user and agree to our Community Guidelines. You will be re-directed back to this page where you will see comments updating in real-time and have the ability to recommend comments to other users.

**Comments**

[Subscribe to comments](#)

There are currently no comments.

**See other News & Comment articles from Nature**

Nature ISSN 0028-0836 EISSN 1476-4687