


Home → Medical Encyclopedia → Fetal alcohol syndrome

Fetal alcohol syndrome



Overview

Infants born to mothers who drink, even in modest amounts, may be born with fetal alcohol syndrome, which often consists of growth retardation, unusual facial features, and intellectual disability.

Review Date 7/16/2024


Updated by: Neil K. Kaneshiro, MD, MHA, Clinical Professor of Pediatrics, University of Washington School of Medicine, Seattle, WA. Also reviewed by David C. Dugdale, MD, Medical Director, Brenda Conaway, Editorial Director, and the A.D.A.M. Editorial team.

[Learn how to cite this page](#)

Was this page helpful?

Yes

No



CERTIFIED
Health Content
Provider
06/01/2028

A.D.A.M., Inc. is accredited by [URAC](#), for Health Content Provider ([www.urac.org](#)). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services.

Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#), and [privacy policy](#).

The information provided herein should not be used during any medical emergency or for the diagnosis or treatment of any medical condition. A licensed medical professional should be consulted for diagnosis and treatment of any and all medical conditions. Links to other sites are provided for information only – they do not constitute endorsements of those other sites. No warranty of any kind, either expressed or implied, is made as to the accuracy, reliability, timeliness, or correctness of any translations made by a third-party service of the information provided herein into any other language. © 1997-2025 A.D.A.M., a business unit of Ebix, Inc. Any duplication or distribution of the information contained herein is strictly prohibited.

